I'm not a robot



TaxonomyThe Blue-spotted Tilapia (Oreochromis leucostictus) is a captivating freshwater fish that has fascinated aquatic enthusiasts and researchers alike. This remarkable species not only mesmerizes with its beauty but also plays a crucial role in its ecosystem. Lets dive into the world of the Blue-spotted Tilapia and explore its unique traits, behaviors, and importance to our environment. Overview The Blue-spotted Tilapia is a species of cichlid that is native to the freshwater lakes and rivers of Africa and parts of the Middle East. Known for its vibrant colors and adaptability, this fish thrives in various aquatic environments. While it holds significant ecological value, the Blue-spotted Tilapia is also popular among aquarists and fish farmers due to its hardiness and nutritional attributes. EtymologyThe scientific name Oreochromis, derives from Greek, meaning mountain color, indicative of their colorful appearances often found in rocky terrains, while the species name leucostictus comes from the Greek words leucos, meaning white, and stictus, meaning spotted, referring to the distinctive blue spots that allows it to navigate through various water currents. Typically, they grow to a length of about 25 to 30 cm (around 10 to 12 inches). Their base color is a shimmering silvery hue, which is beautifully accented by striking blue spots and an assortment of dark markings. These characteristics not only make them visually appealing but also contribute to their identity among cichlids. Identifiable TraitsColoration: The Blue-spotted Tilapia boasts a vibrant silvery body with prominent blue spots and often features darker vertical bars. Dorsal Fin: The dorsal fin is elongated, often extending along the back, characterized by distinct rays and a beautiful gradient of color. Body Shape: They have a typical cichlid shape laterally compressed with a somewhat deep body ideal for maneuvering. Mouth Structure: The mouth is small but well-adapted for grazing algae, which reflects their herbivorous diet. Ecological Significance The Blue-spotted Tilapia plays a pivotal role in its ecosystem. As a herbivore, it primarily feeds on algae and detritus, facilitating the control of algal blooms in freshwater habitats. This feeding behavior helps maintain water quality, which is vital for the overall health of aquatic ecosystems. Moreover, they are a key food source for various predators, thus supporting the food web.Location/RegionOriginally from the tropical and subtropical waters of Africa, the Blue-spotted Tilapia has been introduced in various parts of the world, including Asia and South America. They thrive in freshwater environments, particularly in river valleys, lakes, and wetlands. Their adaptability to different water conditions has enabled them to inhabit both warm and moderate temperatures, making them a resilient species. Social BehaviorBlue-spotted Tilapia exhibit fascinating social dynamics. They are known for forming loose schools, particularly in their juvenile stages, where they often congregate for protection against predators. As they mature, they may become territorial, especially among males vying for mating rights. Mating Practices Tilapia are quite elaborate. During the breeding season, males establish territories and perform courtship displays to attract females. They often create distinct nesting sites in sandy substrates, which they guard zealously. Females lay their eggs in these nests, where both parents typically take an active role in protecting their brood until the fry are free-swimming. This parental care is a remarkable aspect of their reproductive behavior, showcasing the evolutionary strategies they employ to ensure the survival of their offspring. Natural Predators while the Blue-spotted Tilapia is an adept survivor, it is not without its natural enemies. Predatory fish species, such as larger cichlids and catfish, also inhabit the same waters and may pose a threat to young or unsuspecting adults. Birds of prey such as herons and kingfishers also prey on this fish, emphasizing the balance of predator-prey dynamics in their ecosystem. Conservation Status Currently, the Blue-spotted Tilapia is classified as Least Concern according to the IUCN Red List. Their adaptability to various environments and their extensive distribution support their stable population status. However, this does not exempt them from potential threats such as habitat destruction, pollution, and overfishing, particularly in regions where they are heavily exploited for commercial purposes. Human ImpactHuman activities have significantly impacted the habitats of Blue-spotted Tilapia. Their introduction into non-native environments for aquaculture has led to concerns about potential ecological imbalances. However, responsible fishing practices and sustainable aquaculture can mitigate negative impacts on their populations. Educating communities about the importance of maintaining healthy aquatic ecosystems is crucial for the preservation of this species. Interesting FactsThe Blue-spotted Tilapia has been known to change its dietary habits based on available food sources, showcasing their adaptability. They communicate through various vocalizations and body movements, indicating a sophisticated social behavior among cichlids. This species is a staple food fish in many regions, providing essential nutrition to local communities. Blue-spotted Tilapia (Oreochromis leucostictus) is an impressive fish with a vibrant presence in both aquatic ecosystems and human societies. Their ecological role, coupled with their captivating beauty, makes them an essential species worth protecting and studying. From their unique mating practices to their importance in sustaining freshwater habitats, these fish certainly have a tale to tell! So, the next time you encounter a Blue-spotted Tilapia, take moment to appreciate the wondersBlue-spotted Tilapia (Oreochromis leucostictus) Name: Oreochromis leucostictus is a species of cichlid originally native to the Democratic Republic of the Congo and Uganda. It has now been introduced widely in central Africa and is accused of having a negative ecological impact in some areas where introduced. This species can reach a length of 36.3 centimetres (14.3in) SL. This species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments aimed at farming this species can also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be found in the aquarium trade and experiments are also be approximated at a supplication trade and experiments are also be approximated at a supplication trade are also be approximated at a supplication trade ar mossambicus) were illegally introduced into Australian waterways during the 1970s. Both species have become well established in the relatively short time since their introduction. At first they were probably released as unwanted aquarium fish, and Mozambique tilapia are now established in catchments in tropical and subtropical Australia while black mangrove cichlids are mostly restricted to northeast Queensland. Both tilapia species are declared invasive pests in most Australian states. The rapid rate at which both species are spreading, together with native fishes and can dominate local fish communitiesImpacts of Mozambique tilapia reported from overseas include:declines and localised extinctions of native fish species in central America, Sri Lanka, India, Mexico and Venezuelathe collapse of milkfish aquaculture (Chanos chanos) on the Pacific island of Nauru, due to tilapia overcrowding ponds and aggressively out-competing milkfish for food and spacehabitat damage through nest-building activities, including increased bank instability. Reported impacts of black mangrove cichlids in the United States include: aggressive behaviour towards native fishes (eg Sunfish Lepomis spp.)declines in abundance of other native and exotic fishes outnumbering of local fish communities, in some cases making up more than half of the total fish biomassthe potential to reduce the quality of spawning habitat available for native species. Tilapias aggressively compete with native fishes and can dominate local fish communitiesIn Australia, recent detailed studies of the biology of both species have identified potential impacts including activities disruption of spawning in native fishes 1. Environmental impacts Competition: Both species of tilapia introduced to Australia are highly aggressive when breeding and actively defend their territories and young from potential predators. Aggressive behaviour towards Australian native fish has been well documented, with tank trials showing the mere presence of Mozambique tilapia significantly disrupts the breeding success of the Australian eastern rainbow fish1. Mozambique tilapia also have the ability to limit their physical size (stunt) and mature at an early age2. This means they can build up their population faster than native fish that cant reproduce at the same rate. This rapid reproduce at the same rate. fishes and become the most dominant fish species where they are introduced. Habitat degradation: Tilapia can cause changes and damage to aquatic habitats. Black mangrove cichlids and Mozambique tilapia both dig out hollows or nests in the river bed during breeding3. When there are a lot of tilapia, large amounts of nest-building activities can cause damage to the bottom of waterways and harm aquatic plants. The damage caused by tilapia might not be as obvious as the damage caused to river bottoms by the feeding activities of introduced carp, but scientific research has shown that Mozambique tilapia can reduce the population density of some Australian native aquatic plants4. Disease: There is potential for introduced fish to bring new types of pathogens and parasites with them. To date, there have been no confirmed cases of either species of tilapia introducing exotic diseases or parasites to Australian waters. However, its possible that the non-native external parasite Trichodina heterodentata was introduced to Australia with tilapia in particular) cultured in the Philippines. Economic impacts Water quality: The introduction of tilapia can decrease water quality in warm-water lakes and reservoirs. Evidence of this impact occurring overseas is from a different tilapia than those we have here6; however, Australian water managers might be reluctant to draw down dams to low levels if reservoirs contain large quantities of tilapia, for fear of mass tilapia deaths fouling the water. Other events resulting in large numbers of fish dying, such as cold weather (which has happened in southeast Queensland7), also have the potential to foul domestic water supplies. Such events could have substantial economic flow-on effects if towns/ cities are forced to buy water from alternative sources, and/or spend money on reducing the number of tilapia in the waterbody. Commercial and recreational fisheries: The effect of introduced tilapia on fisheries resources has not been studied in detail in Australia. Competition for food, aggressive behaviour, habitat degradation and disease transfer are all potential impacts of tilapia on recreational and commercial fisheries. Despite this, in some countries Mozambique tilapia and similar species have been deliberately introduced to try and enhance local fisheries. One study on fish catch rates in a reservoir in Brazil where the closely related Nile tilapia (Oreochromis niloticus) had been introduced showed that fisheries were not actually enhanced. The catch rates of local fishers over a 30- year period were significantly reduced for commercially important native species, but increased for introduced Nile tilapia giving no overall increase in the productivity of the fishery as a whole. In Australia, commercial fisheries are mainly coastal and estuarine, but some of the target species such as barramundi need freshwater systems during parts of their lifecycle. So, any adverse impacts on freshwater systems caused by pest fish such as tilapia could impact on the productivity of associated fisheries. Similarly, invasive tilapia populations have the potential to affect recreational fisheries by increasing the amount of trash fish caught by anglers. This might have flow-on economic impacts on related industries such as tourism, especially if recreational fisheries are an important part of the local economy. Social impacts: People are less likely to use and enjoy public reservoirs and waterways if they contain too many pest species such as tilapia. For example, water bodies fouled by nest-building activities, the presence of dead fish, or even changes in what types of fish are caught, can all change the way the community perceives and uses the resource. Further information: Doup RG, Schaffer J, Knott MJ and Burrows D (2009). How might an exotic fish disrupt success in a sympatric native species? Marine and Freshwater Research 60:379383. Russell DJ, Thuesen PA and Thomson FE (2010). Development of Management Strategies for Control of Feral Tilapia Populations in Australia. Queensland Department of Employment, Economic Development and Innovation (DEEDI), Brisbane, Queensland. Doup RG and Burrows DW (2008). Thirty years later, should we be more concerned for the ongoing invasion of Mozambique tilapia in Australia? Pacific Conservation Biology 14:235238. Doup RG, Knott MJ, Schaffer J, Burrows D and Lymbery AJ (2010). Experimental herbivory of native Australian macrophytes by the introduced Mozambique tilapia Oreochromis mossambicus. Austral Ecology 35:2430.Dove ADM and ODonoghue PJ (2005). Trichodinidae) from native and exotic Australian freshwater fishes. Acta Protozoologica 44:5160.Starling F, Lazzaro X, Cavalcanti C and Moreira R (2002). Contribution of omnivorous tilapia to eutrophication of a shallow tropical reservoir: evidence from a fish kill. Freshwater Biology 47:24432452. Sawynok W, Parsons W and Platten P (2009). Winter 2007 Fish Death Events: Impact on Stocked Fish. Infofish Services, Rockhampton, Queensland. Attayde JL, Brasil J and Menescal RA (2011). Impacts of introducing Nile tilapia on the fisheries of a tropical reservoir inAgnse, J.F. and Gourne, B.P. (1998) Natural hybridization between Tilapia zillii (Gervais, 1848) and T. guineensis (Bleeker, 1962) in Ivory Coast, in Abstracts from the Symposium on African Fishes and Fisheries: Diversity and Utilization, Grahamstown, South Africa, 1998, p. 14. Google Scholar Allanson B.R. (1979) Lake Sibaya, in Lake Sibaya, in Lake Sibaya, in Lake Sibaya. Mongr. Biol. 36, Dr W. Junk, The Hague, pp. 313328. Google Scholar Arthington, A.H. and Bluhdorn, D.R. (1996) The effects of species introductions resulting from aquaculture operations, in Aquaculture and Water Resource Management (eds D.J. Baird, M.C.M. Beveridge, L.A. Kelly and J.F. Muir), Blackwell, Oxford, pp. 114139. Google Scholar Bailey, R.G. (1966) The dam fisheries of Tanzania. E. Afric. Agricult. Forest. J. 32(1), 115. Google Scholar Bailey, R.G. (1988) Fish and fisheries, in The Jonglei Canal (eds P. Howell, M. Lock and S. Cobb), Cambridge University Press, Cambridge, pp. 328349. Google Scholar Bailey, R.G. (1996) Changes in the fish and fisheries ecology of a large man-made lake in Tanzania, 1965-94. Fish. Manage. Ecol. 3, 251260. Article Google Scholar Balirwa, J.S. (1998) Lake Victoria Wetlands and the Ecology of the Nile tilapia Oreochromis niloticus L., PhD dissertation, Agricultural University, Wageningen, Balkema, Rotterdam, the Netherlands. Google Scholar Balogun, J.K. and Ibeun, M.O. (1995) Additional Information on Fish Stocks and Fisheries of Lake Kaini (Nigeria). CIFA Technical Paper, 30, FAO, Rome, pp. 1940. Google Scholar Baluyut, E.A. (ed.) (1992) Papers Contributed to the Workshop on Tilapia in Capture and Culture-Enhanced Fisheries in the Indo-Pacific Fishery Commission Countries, Bogor, Indonesia, June 1991. FAO, Rome. Google Scholar Beauchamp, R.S.A. (1964) The Rift Valley lakes of Africa. Verh. Int. Verein. Limnol. 15, 9199 Google Scholar Bootsma, H.A., Hecky, R.E., Hesslein, R.H. and Turner. G.F. (1996) Food partitioning among Lake Malawi nearshore fishes as revealed by stable isotope analyses. Ecology 77, 12861290. Article Google Scholar Bowen, S.H. (1979) A nutritional constraint in detritivory by fishes: the stunted population of Sarotherodon mossambicus in Lake Sibaya, South Africa. Ecol. Monogr. 49, 1731. Article Google Scholar Bowen, S.H. (1980) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela. Science 207, 12161218. Article PubMed CAS Google Scholar Bowen, S.H. (1984) Detrital non-protein amino acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela and Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth acids are the key to rapid growth of Tilapia in Lake Valencia, Venezuela acids are the key to rapid growth acids are the key to Fishes (eds C. Leveque, M.N. Bruton and G.W. Ssentongo), Travaux et Documents, 216, ORSTOM, Paris, pp. 243247. Google Scholar Braimah, L.I. (1995) Recent developments in the fisheries of Volta Lake (Ghana), CIFA Technical Paper, 30, FAO, Rome, pp. 111134. Google Scholar Bruton, M.N. (1979) The fishes of Lake Sibaya, in Lake Sibaya (ed B.R. Allanson), Mongr. Biol. 36, Dr W. Junk, The Hague, pp. 162245. Google Scholar Burgis, M.J. and Dunn, I.G. (1978) Production in three contrasting ecosystems, in Freshwater Fish Productions into the Sepik River, Papua New (East Africa). IHP-IV Project M-5.1 SC-95/WS/16 UNESCO, Paris. Google Scholar Duponchelle, F. and Panfili, J. (1998) Variations in age and size at maturity of female Nile tilapia Oreochromis niloticus, populations from man-made lakes of Cote dIvoire. Env. Biol. Fishes 52, 453465. Article Google Scholar FAO (1993) Fisheries Management in the Southeast Arm of Lake Malawi, the Upper Shire River and Lake Malombe, With Particular Reference to the Fisheries on Chambo (Oreochromis spp). CIFA Technical Paper, 21, FAO, Rome. Google Scholar Fryer, G. (1961) Observations on the biology of the cichlid fish Tilapia variabilis in the northern waters of Lake Victoria (East Africa). Rev. Zool Bot. Afr. 64, 133. Google Scholar Gophen, M., Pollingher, U. Ochumba, P.O.B. and Kaufman, L.S. (1993) Nile perch (Lates niloticus) invasion in Lake Victoria (East Africa). Verh. Internat. Verein. Limnol. 25, 856859. Google Scholar Gourne, G., Teugels, G.G., Hugueny, B. and Thys van den Audenaerde, D.F.E. (1998) Evaluation of the icthyological diversity of a West African basin after the construction of a dam (Bia River, Ivory Coast and Ghana), in Abstracts, Symposium on African Fishes and Fisheries, Grahamstown, South Africa, 1998, p. 42. Google Scholar Greenwood, P.H.G. and Lund, J.W.G. (1976) A discussion on the biology of an equatorial lake: Lake George, Uganda. Proc. Roy. Soc Lond. 184B, 227346. Google Scholar Guerrero, III, R.D. (1999) Impacts of tilapia introductions on the endemic fishes in some Philippine lakes and Reservoirs, in Fish and Fisheries of Lakes and Reservoirs, in Fish and Fisheries of Lakes and Reservoirs in SE Asia and Africa (eds W.L.T. van Densen and M.I. Morris), Westbury Academic and Scientific Publishing, Otley, UK, pp. 151157. Google Bangor. Google Scholar Hecky, R.E. (1993) The eutrophication of Lake Victoria. Kilham Memorial Lecture, 25th Congress of SIL, Verh. Int. Verein. Limnol. 25, 3948.CAS Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Hydrobiologia 144, 243250.Article Google Scholar Hickley, P. and Bailey, R.G. (1987) Fish communities in the eastern seasonal-floodplain of the Sudd, Southern Sudan. Scholar Holden, M.J. (1963) The Populations of Fish in Dry Season Pools of the River Sokoto. Colonial Office, London. Google Scholar Hopson, T. et al. (eds) (1982) Lake Turkana. A Report on the Findings of the Lake Turkana. Project 1972-1975, ODA, London. Google Scholar Iles, T.D. (1973) Dwarfing or stunting in the genus Tilapia (Cichlidae) a possibly unique recruitment mechanism. Rapp. Reun. Cons. Perm. Int. Explor. Mer. 164, 247254. Google Scholar Ita, E.O. (1984) Lake Kainji (Nigeria), in Status of African Reservoir Fisheries (eds J.M. Kapetsky and T. Petr), CIFA Technical Paper, 10, FAO, Rome, pp. 43103. Google Scholar Johnson, T.C. Scholz, CA., Talbot, M.R., Kelts, K., Ricketts, R.D., Ngobi, G., Beuning, K., Ssemmanda, I. and McGill, J.W. (1996) Late Pleistocene desiccation of Lake Victoria and rapid evolution of cichlid fishes. Science 273, 10911093. Article PubMed CAS Google Scholar Juarez-Palacios J.R. and Olmos-Tomassini, M.E. (1992) Tilapia in Capture and Cultureenhanced Fisheries in Latin America. FAO Fish, Technical Report 458, Suppl., pp. 244273. Google Scholar Kaufman, L. (1992) Catastropic change in species-rich freshwater ecosystem of Lake Turkana: an ever-changing mixed environment. Mitt Verein. Internat. Limnol. 23, 2535. Google Scholar Kolding, J. (1993) Population dynamics and life-history styles of Nile tilapia, Oreochromis niloticus, in Fergusons Gulf, Lake Turkana, Kenya, Env. Biol. Fishes 37, 2546. Article Google Scholar Kolding, J. (1995) Changes in species composition and abundance of fish populations in L. Turkana, Kenya, in The Impact of Species Changes in African Lakes (eds T.J. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 335363. Chapter Google Scholar Lauzanne, L. (1972) Regimes alimentaires principales especs de poissons de larchipel oriental du lac Tchad. Verh. Internat. Verein. Limnol. 18, 636646. Google Scholar Lehman, J.T. (d.) (1998) Environmental Change and Response in East African Lakes, Kluwer Academic Press, London. Google Scholar Linn, I. J. and K.L.I. Campbell. (1986) Cormorants and Fisheries. A Report on the Biology of the White-breasted Cormorant Phalacrocorax carbo as it Affects the Commercial Fisheries of Lake Malawi. Overseas Development Administration, London. Google Scholar Lowe, R.H. (1952) Report on the Tilapia and Other Fish and Fisheries of Lake Nyasa 1945-47, Colonial Office, London, Fishery Publ. 1(2), 1126. Google Scholar Lowe, R.H. (1955a) New species of Tilapia (Pisces, Cichlidae) from Lake Jipe and the Pangani River, East Africa. Bull. Brit. Mus. (Nat. Hist.) 2, 349368. Google Scholar Lowe (McConnell), R.H. (1955b) The fecundity oiTilapia species. East Afr. Agric. J. 11, 4552. Google Scholar Lowe (McConnell), R.H. (1956a) Observations on the biology of Tilapia (Pisces-Cichlidae) in Lake Victoria, East Afr. Fish. Res. Org, Suppl. Pub. 1, 55, 353373. Google Scholar Lowe (McConnell), R.H. (1958) Observations on the biology of Tilapia nilotica Linne in East African waters. Rev. Zool. Bot. Afr. 57, 129170. Google Scholar Lowe (McConnell), R.H. (1959) Breeding behaviour patterns and ecological differences between Tilapia species and their significance for evolution within the genus Tilapia (Pisces: Cichlidae). Proc. Zool. Soc. (Lond) 132, 130. Google Scholar Lowe-McConnell, R.H. (1982) Tilapias in fish communities, in The Biology and Culture of Tilapias (1CLARM, Manila, pp. 83114. Google Scholar Lowe-McConnell, R.H. (1987) Ecological studies of Tropical Fish Communities, Cambridge University Press, Cambridge Book Google Scholar Man, H.S.H. and Hodgkiss, I.J. (1977) Studies on the ichthyofauna in Plover Cove Reservoir, Hong Kong: feeding and food relations. J. Fish Biol. 11, 113. Article Google Scholar Mattson, N.S. and Kaunda, E.K.W.H. (1997) Population dynamics of Oreochromis shiranus in two small water bodies in Malawi. J. Fish Biol. 50, 592607. Google Scholar Moreau, J., Arrignon, J. and Jubb. R.A. (1988) Introduction of foreign fishes in African inland waters. Suitability and problems, in Biology and Ecology of African Freshwater Fishes (eds C. Leveque, M.N. Bruton and G.W. Ssentongo), Travaux et Documents No 216, ORSTOM, Paris, pp. 395425. Google Scholar Moreau, J., Christensen, V. and Pauly, D. (1993) A trophic ecosystem model of Lake George, Uganda, in Ecotrophic Models of Aquatic Ecosystems (ICLARM Conference Proceedings, Vol. 26) (eds V. Christensen and D. Pauly), ICLARM, Manila, pp, 124129. Google Scholar Moriarty D.J.W., Darlington, J.P.E.C., Dunn, I.G., Moriarty, CM and Tevlin, M.P. (1973) Feeding and grazing in Lake George, Uganda. Proc. R. Soc. Lond. 184B, 299319.Article Google Scholar Muchiri, S.M., Hart, P.J.B. and Harper, D.M. (1995) The persistence of two introduced tilapia species in Lake Naivasha, Kenya, in the face of environmental variability and fishing pressure, in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 291319. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and Kaufman, L. (1995) A note on recent advances in the genetic characterization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and W. Alexanderization of Tilapiine stocks in Lake Victoria Region. Afr. J. Trop. Hydrobiol. Fish. 6, 5153. Google Scholar Mwanja, W. and W. Alexander Noakes, D.L.G. and Balon, E.K. (1982) Life histories of tilapias: an evolutionary perspective, in The Biology and Culture of Tilapias (ICLARM, Manila, pp. 6180. Google Scholar Ogutu-Ohwayo, R. (1992) The purpose, costs and benefits of fish introductions; with specific reference to the Great Lakes of Africa. Mitt Internat. Verein. Limnol. 23, 3744. Google Scholar Ogutu-Ohwayo, R. 1995. Diversity and stability of fish stocks in Lake Victoria, Kyoga and Nabugabo after the establishment of introduced species, in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 5981. Chapter Google Scholar Palsson, O.K., Bulirani, A. and Banda, M. (1998) A review of the population dynamics of chambo Oreochromis spp. (Cichlidae) in Lakes Malawi and Malombe, in Abstracts from the Symposium on African Fishes and Fisheries: Diversity and Utilization, Grahamstown, South Africa, 1998, p. 158. Google Scholar Pauly, D. (1976) The biology, fishery and potential for aquaculture of Tilapia melanotheron in a small West African lagoon. Aquaculture 7, 3349. Article Google Scholar Pauly, D. (1984) A mechanism for the juvenile-to-adult transition in fishes. J. Cons. Int. Expl. Mer. 41, 280284. Google Scholar Pet, J.S. and Piet, G.J. (1993) The consequences of habitat occupation and habitat overlap of the introduced tilapia O. mossambicus and indigenous fish species in resource management in a Sri Lankan reservoir. J. Fish Biol. 43(Suppl. A), 193208. Article Google Scholar Petr, T. (1967) Fish population changes in the Volta Lake in Ghana during its first sixteen months. Hydrobiologia 30, 193220. Article Google Scholar Philippart, J.-CL, and Ruwet, J.-Cl. (1982) Ecology and distribution of Tilapias, in The Biology and Culture of Tilapias (ICLARM, Manila, pp. 1559. Google Scholar Piet, F.J., Pet, J.S. and Guruge, W.A.H.P. (1994) Niche partitioning of indigenous riverine carps and the exotic lacustrine tilapia (Oreochromis mossambicus) in a reservoir in SE Sri Lanka. Verh. Internat. Verein. Limnol. 25, 21832187. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Changes in African Lakes, Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and Hart, P.J.B. (eds) (1995) The Impact of Species Chapman and Hall, London. Google Scholar Pitcher, T.J. and T Biologie et Exploitation des Poissons du Lac Ihema (bassin Akagera), Rwanda, Presses Universitaires de Namur, Belgium. Google Scholar Schiemer, F. (1996) Significance of filter-feeding fish in tropical waters, in Perspectives in Tropical Limnology (eds F. Schiemer and K.T. Bolland), SPB Academic by, Amsterdam, pp. 6575. Google Scholar Schliewen, U.K., Tauz, D. and Paabo, S. (1994) Sympatric speciation suggested by monophyly of crater lake cichlids. Nature (Lond.) 368, 629632. Article CAS Google Scholar Seegers, L. (1996) The Fishes of the Lake Rukwa Drainage. Ann. Mus. r. Afr. Cent. Sci. Zool. 278, 407. Google Scholar Stiassny, and Tichy, H. (1999) The Oreochromis alcalicus flock from Lakes Natron and Magadi, Tanzania and Kenya, with descriptions of two new species. Ichthyol. Explor. Freshwaters 10, 97146. Google Scholar Stiassny, M., Schliewen, U.K. and Dominey, W.J. (1992) A new species flock of cichlid fishes from Lake Bermin, Cameroon with a description of eight new species of Tilapia (Labroidei: Cichlidae). Ichth. Expl. Freshwat. 3, 311346. Google Scholar Trewavas, E. (1983) Tilapiine Fishes of the Genera Sarotherodon, Oreochromis and Danakilia, British Museum (Natural History) London. Google Scholar Trewavas, E., Green, J. and Corbet, S.A. (1972) Ecological studies on crater lakes in West Cameroon. Fishes of Barombi Mbo. J. Zool. (Lond). 167, 4195. Article Google Scholar Turber, G.W. (1995) Management, conservation and species changes of exploited fish stocks in Lake Malawi, in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 365395. Chapter Google Scholar Tweddle, D., Turner, G.W. and Seisay, M.B.D. (1995) Changes in species composition and abundance as a consequence of fishing in Lake Malombe, Malawi, in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 413424. Chapter Google Scholar Twongo, T. (1995) Impact of fish species introductions on the tilapias of Lakes Victoria and Kyoga, in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 4556. Chapter Google Scholar Vareschi, E. (1979) The ecology of Lake Nakuru (Kenya) II Biomass and spatial distribution of fish (T. grahami). Oecologia (Berlin) 37, 321335. Google Scholar Welcomme, R.L. (1964a) Notes on the present distribution and habits of the non-endemic species of Tilapia which have been introduced into Lake Victoria. E. Afr. Fish. Res. Org. Ann. Kept. 1962/63, 3639. Google Scholar Welcomme, R.L. (1964) The habitats and habitat preferences of the young of the Lake Victoria Tilapia (Pisces, Cichlidae). Rev. Zool. Bot. Afr. 70, 128. Google Scholar Welcomme, R. L. (1966) Recent changes in the stocks of Tilapia in L. Victoria. Nature 212, 5254. Article Google Scholar Welcomme, R.L. (1967b) The relationship between fecundity and fertility in the mouth-brooding cichlid fish Tilapia leucosticta. J. Zool. (London) 151, 453468. Google Scholar Welcomme, R.L. (1968) Observations on the biology of the introduced species of tilapia in Lake Victoria. Rev. Zool Bot. Afr. 76, 249279. Google Scholar Welcomme, R.L. (1970) Studies on the effects of abnormally high waterlevels on the ecology of fish in certain shallow regions of L. Victoria. J. Zool. (London) 160, 405436. Article Google Scholar Welcomme, R.L. (1988) International Introductions of Inland Aquatic Species. FAO Fish. Technical Paper, 294, FAO, Rome. Google Scholar Witte, F., Goldschmidt, T. and Wanink, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in The Impacts of Species Changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 83135. Chapter Google Scholar Worthington, E.Brackette, F., Goldschmidt, T. and Wanink, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 83135. Chapter Google Scholar Worthington, E.Brackette, F., Goldschmidt, T. and Wanink, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 83135. Chapter Google Scholar Worthington, E.Brackette, F., Goldschmidt, T. and Wanink, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 83135. Chapter Google Scholar Worthington, E.Brackette, F., Goldschmidt, F. and Wanink, J.H. (1995) Dynamics of the haplochromine cichlid fauna and other ecological changes in African Lakes (eds T. Pitcher and P.J.B. Hart), Chapman and Hall, London, pp. 83135. Chapter Google Scholar Worthington, E.Brackette, F. and (1937) On the evolution of fish in the Great Lakes of Africa. Int. Rev. gesamt. Hydrobiol. Hydrog. 35, 304317.Article Google Scholar Worthington, E.B. and Ricardo, C.K. (1936) Scientific Results of the Cambridge Expedition to the East African Lakes, 1930-1. No 15. The fish of Lake Baringo. J. Linn. Soc. Zool. 39, 353389. Article Google Scholar Just over 40 years after introduction into the wild near Brisbane, Mozambique tilapia (Oreochromis mossambicus) are now widespread throughout coastal catchments in Queensland and crossed the border into the New South Wales Tweed catchment for the first time in 2014. In 2022, Australia reported the greatest number of new Tilapia invasions globally, with new populations in Queensland and New South Wales. Tilapia are expected to adversely impact 19 of the 46 species of native fish in the Murray-Darling Basin; either through direct predation, introduction of diseases, competition for food and habitat, and/or habitat destruction. Impacts on water quality through increased nutrient cycling, winter die-offs, loss of aquatic vegetation and nest construction are also predicted. High riskSeveral Tilapia infested catchments border the north-eastern section of the Murray-Darling Basin. In some locations, such as the Lockyer catchment, Tilapia have been observed within 10 km of the Murray-Darling Basin. Human assisted movement (intentional or accidental) of Tilapia is the biggest threat and most likely pathway for a Murray-Darling Basin incursion. Mozambique tilapia. Photo credit: Townsville BulletinThe Condamine and Border Rivers catchments in the north-east of the Murray-Darling Basin are the highest risk areas for a Tilapia incursion due to their close proximity to infested Queensland catchments. However, with human assisted movement Tilapia could turn up anywhere. Basin reportsIn 2017, a suspected Tilapia sighting was reported in the Barwon River in western New South Wales. Fortunately, subsequent monitoring by New South Wales Department of Primary Industries failed to detect the presence of Tilapia in the area, and officers concluded that the report was most likely a misidentification. Alarmingly, this misidentified Tilapia catch was not reported to authorities for nearly three months. In a system like the Barwon Darling, eradication or even control of Tilapia three months after introduction would have been impossible in all but the lowest of flow conditions (i.e. unless the system ceased to flow). Advisory materials have been reissued to this and other areas of New South Wales to increase efforts for rapid reporting of future suspect reports of Tilapia. Water temperatures across large areas of the northern Murray-Darling Basin are suitable for Tilapia breeding for approximately six months of the year (warmer months). During a six-month breeding season, females would be able to brood at least five times, significantly increasing the introduced population and hindering eradication efforts further. Tilapia can also effectively move upstream and downstream. In the Burdekin River, northern Queensland, Tilapia were recorded spreading 3,000 km just four years after introduction. Adult: Mozambique Tilapia have pale olive to silver-grey bodies, with a long continuous dorsal (top) fin that ends in a sharp point, and can grow to more than 36 cmControl optionsTilapia eradication management options are largely limited to the use of rotenone, which has strict conditions for its use under an Australian Pesticides and Veterinary Medicines Authority Permit. Rotenone is a naturally occurring plant derived chemical, which inhibits cellular respiration, and has seen unintended consequences for native fish Lethal concentrations of rotenone are non-selective to fish, killing all fish in the treated area. As such, its use in pest fish control must make both ecological and economic sense and is largely limited to use in small waterways and storages that can be isolated from surrounding waters with regulating structures or bunds. Other control tools for Tilapia include electrofishing and chemical applications, however, education is the most important tool to prevent additional incursions. Incursions outside the Murray-Darling BasinIn 2017, another report was made of a Spotted tilapia (Tilapia mariae) sighting in the Walsh River, part of the Mitchell River Catchment in Far North Queensland. The Walsh River (previously thought to be Tilapia free) is a western flowing waterway that drains into the Gulf of Carpentaria where productive commercial and recreational prawn and Barramundi fisheries exist. Biosecurity Queensland have confirmed Spotted tilapia are present in the Walsh River. Media coverage of this news generated further independent community reports of Tilapia presence in the nearby Mitchell River for two years. Despite extensive electrofishing and netting at the reported locations in the Mitchell River, Biosecurity Queensland has failed to validate those reports. The extent of the Walsh River Tilapia infestation is still being investigated. With the wet season in far north Queensland underway, Tilapia will likely spread further throughout the catchment towards the Gulf. The rugged remoteness and presence of crocodiles in many areas complicates the management response even further. In 2008, the Endeavour River Catchment in North Queensland became infested after a private landholder deliberately stocked property dams with Mozambique tilapia. Subsequent flooding events saw the Tilapia spread to nearby Jensens Crossing on the Endeavour River. The catchment is now infested and eradication not possible. Juvenile Stunded Mossambicus Community awareness The vast and largely remote expanse of the Murray-Darling Basin makes monitoring for potential Tilapia incursions extremely difficult. River users, recreational fishers, landholders, local government and natural resource management agencies play a critical role in both keeping Murray-Darling Basin free of Tilapia, as well as being the eyes and ears on the ground. Social and economic impacts of a Tilapia incursion are also a reality. Costs associated with eradication and management, increased water treatment, winter die-off clean ups, and losses in recreational fishing trade and tourism may all potentially impact Murray-Darling Basin communities. Recent examples of new Tilapia incursions in far north Queensland and northern coastal New South Wales are a significant concern. Despite education and awareness programs and strict regulatory controls, Tilapia are still being moved between catchments. Without doubt the best form of Tilapia control in the Murray-Darling Basin is prevention. How you can helpLearn how to correctly identify Tilapia are still being moved between catchments. Without doubt the best form of Tilapia control in the Murray-Darling Basin is prevention. How you can helpLearn how to correctly identify Tilapia are still being moved between catchments. Without doubt the best form of Tilapia control in the Murray-Darling Basin is prevention. of Tilapia to the appropriate authoritiesDont throw caught Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings, illegal movement or stocking as soon as possibleThe Keep Tilapia as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive)Report Tilapia catches and sightings as bait (dead or alive) funding from the Murray-Darling Basin Authority, Tilapia identification: Department of Agriculture and FisheriesReport by phone: 13 25 23Report online here. Report by email: pestfish@daf.gld.gov.auNew South WalesInformation and identification: Department of Primary IndustriesReport by phone (02) 4916 3877Report online here.Report by email: aquatic.pests@dpi.nsw.gov.auReferencesHutchison M., Sarac Z. and Norris A. (2011) Mozambique Tilapia, the Potential for Mozambique Tilapia (Oreochromis Mossambicus) to Invade the Murray-Darling Basin and the Likely Impacts: a Review of Existing Information Murray-Darling Basin Authority, Commonwealth of Australia. The State of Queensland (2011) Stop the Spread Tilapia Training Package. The State of Queensland, Department of Employment, Economic Development and Innovation, Brisbane. Tilapia are members of the Cichlid family and are one of many invasive noxious fish species that have spread across many parts of the world causing irreversible damage. Tilapia were introduced into Australia in the 1960s and now dominate most major Queensland catchments. There are 2 main species within Australian waterways the Mozambique tilapia (Oreochromis mossambicus) and the spotted tilapia (Tilapia mariae). The Murray-Darling Basin is now the frontline for these fish. Unfortunately these fish are easily spread by people dumping them as bait and from natural and human water transfers. Once these fish establish themselves they are difficult, if not impossible, to eradicate. Native fish species are outcompeted and attacked by tilapia species which are highly aggressive and may dominate waterways causing a detrimental shift in riverine and wetland ecosystem function. The biological characteristics that allow them to dominate waterways causing a detrimental shift in riverine and wetland ecosystem function. The biological characteristics that allow them to dominate waterways causing a detrimental shift in riverine and wetland ecosystem function. many native fish cannot. Female Tilapia carry and aggressively defend fertilised eggs and young in their mouth, resulting that energy into reproducing at an earlier age rather than putting that energy into growth. Turbid, warm and polluted waters low in oxygen are no problem for Tilapia as they can survive temperatures from 8 42C and gulp air from the surface if needed. They can also survive well in seawater. Being omnivorous, they consume anything from algae to other fish species, including their own. As tilapia disrupt the function of native aquatic ecosystems they also indirectly cause a decline in water quality. They can disturb sediment and in-stream plant life by grazing and from extensive construction of nests in substrate. How does this relate to bush or riparian regeneration and conservation? The answer is that Tilapia are more of a symptom of degraded riverine and wetland habitat rather than the cause. They generally prefer against Tilapia (and other pest fish) incursions. This can be done by taking a holistic approach to river rehabilitation. For example we can increase the resilience of less competitive native fish habitat by restoring riparian and in-stream vegetation and by re-snagging creek banks (also by leaving fallen trees in waterways). In addition, removal of fish barriers such as weirs, culverts and crossings can help improve resilience of native fish species by allowing passage for feeding and reproduction. Article by Daniel Bloom, waterways officer, Sunshine Coast and Freshwater Research 62: 902917. AbstractCadwallader, P.L., Backhouse, G.N. & Fallu, R. (1980). Occurrence of exotic tropical fish in the cooling pondage of a power station in temperate south-eastern Australia. Australia. Australia. Australia. Freshwater Research 31: 541546. AbstractCanonico, G.C., Arthington, A., McCrary, J.K. & Thieme, M.L. (2005). The effects of introduced tilapias on native diversity. Aguatic Conservation: Marine and Freshwater Ecosystems 15: 463483. Dunz, A.R. & Schliewen, U.K. (2013): Molecular Phylogenetics and Evolution 68(1): 6480. doi: 10.1016/j.ympev.2013.03.015 AbstractMather, P.B. & Arthington, A. H. (1991). Assessment of genetic differentiation among feral Australian tilapia populations. Australian tilapia populations on the introduced tropical cichlid, Tilapia mariae. B.Sc. (Hons) Thesis, Latrobe University, Melbourne.Russell DJ, Thuesen PA & Thomson FE (2009). Development of Management Strategies for Control of Feral Tilapia Populations in Australia. Queensland Department of Employment, Economic Development and Innovation, Brisbane.Webb, A.C. (2007). Status of non-native freshwater

Parts of speech in english pdf free download. Parts of speech pdf in english. Parts of speech in gujarati pdf with examples. Parts of speech in gujarati. Gujarati speech pdf.